#84

NOAA Technical Report NMFS 84

December 1989

Results of Abundance Surveys of Juvenile Atlantic and Gulf Menhaden, *Brevoortia tyrannus* and *B. patronus*

Dean W. Ahrenholz
James F. Guthrie
Charles W. Krouse

LIBRARY

JAN 29 1990

National Marine Fisheries Service
2570 Dole Street
Honolulu, Hawaii

NOAA
Tech.
Rept.
NMFS
84
c.2

U.S. Department of Commerce

NOAA TECHNICAL REPORT NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS; intensive scientific reports on studies of restricted scope; papers on applied fishery problems; technical reports of general interest intended to aid conservation and management; reports that review in considerable detail and at a high technical level certain broad areas of research; and technical papers originating in economics studies and from management investigations. Since this is a formal series, all submitted papers receive peer review and those accepted receive professional editing before publication.

Copies of NOAA Technical Reports NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161. Although the contents have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

48. Widow rockfish: Proceedings of a workshop, Tiburon, California, December 11–12, 1980, by William H. Lenarz and Donald R. Gunderson (editors). January 1987, 57 p.

49. Reproduction, movements, and population dynamics of the southern kingfish, *Menticirrhus americanus*, in the northwestern Gulf of Mexico, by Stephen M. Harding and Mark E. Chittenden, Jr. March 1987, 21 p.

50. Preparation of acetate peels of valves from the ocean quahog, *Arctica islandica*, for age determinations, by John W. Ropes. March 1987, 5 p.

51. Status, biology, and ecology of fur seals: Proceedings of an international workshop, Cambridge, England, 23–27 April 1984, by John P. Croxall and Roger L. Gentry (editors). June 1987, 212 p.

52. Limited access alternatives for the Pacific groundfish fishery, by Daniel D. Huppert (editor). May 1987, 45 p.

53. Ecology of east Florida sea turtles: Proceedings of the Cape Canaveral, Florida, sea turtle workshop, Miami, Florida, February 26–27, 1985, by Wayne N. Witzell (convener and editor). May 1987, 80 p.

54. Proximate and fatty acid composition of 40 southeastern U.S. finfish species, by Janet A. Gooch, Malcolm B. Hale, Thomas Brown, Jr., James C. Bonnet, Cheryl G. Brand, and Lloyd W. Reiger. June 1987, 23 p.

55. Proximate composition, energy, fatty acid, sodium, and cholesterol content of finfish, shellfish, and their products, by Judith Krzynowek and Jenny Murphy. July 1987, 53 p.

56. Some aspects of the ecology of the leatherback turtle *Dermochelys coriacea* at Laguna Jolova, Costa Rica, by Harold F. Hirth and Larry H. Ogren. July 1987, 14 p.

57. Food habits and dietary variability of pelagic nekton off Oregon and Washington, 1979–1984, by Richard D. Brodeur, Harriet V. Lorz, and William G. Pearcy. July 1987, 32 p.

58. Stock assessment of the Gulf menhaden, *Brevoortia patronus*, fishery, by Douglas S. Vaughan. September 1987, 18 p.

59. Atlantic menhaden, *Brevoortia tyrannus*, purse seine fishery, 1972–84, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, William R. Nicholson, Douglas S. Vaughan, Donnie L. Dudley, and Ethel A. Hall. September 1987, 23 p.

60. Gulf menhaden, *Brevoortia patronus*, purse seine fishery, 1974–85, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, Eldon J. Levi, Douglas S. Vaughan, and Ethen A. Hall. December 1987, 8 p.

61. Manual for starch gel electrophoresis: A method for the detection of genetic variation, by Paul B. Aebersold, Gary A. Winans, David J. Teel, George B. Milner, and Fred M. Utter. December 1987, 19 p.

62. Fishery publication index, 1980–85; Technical memoradum index, 1972–85, by Cynthia S. Martin, Shelley E. Arenas, Jacki A. Guffey, and Joni M. Packard. December 1987, 149 p.

63. Stock assessment of the Atlantic menhaden, *Brevoortia tyrannus*, fishery, by Douglas S. Vaughan and Joseph W. Smith. January 1988, 18 p.

64. Illustrated key to penaeoid shrimps of commerce in the Americas, by Isabel Pérez Farfante. April 1988, 32 p.

65. History of whaling in and near North Carolina, by Randall R. Reeves and Edward Mitchell. March 1988, 28 p.

66. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific, by M. James Allen and Gary B. Smith. April 1988, 151 p.

67. Index numbers and productivity measurement in multispecies fisheries: An application to the Pacific coast trawl fleet, by Dale Squires. July 1988, 34 p.

68. Annotated bibliography II of the hard clam *Mercenaria mercenaria*, by J.L. McHugh and Marjorie W. Sumner. September 1988, 59 p.

69. Environmental quality and aquaculture systems: Proceedings of the thirteenth U.S.–Japan meeting on aquaculture, Mie, Japan, October 24–25, 1984, edited by Carl J. Sindermann. October 1988, 50 p.

70. New and innovative advances in biology/engineering with potential for use in aquaculture: Proceedings of the fourteenth U.S.–Japan meeting on aquaculture, Woods Hole, Massachusetts, October 16–17, 1985, edited by Albert K. Sparks. November 1988, 69 p.

71. Greenland turbot *Reinhardtius hippoglossoides* of the eastern Bering Sea and Aleutian Islands region, by Miles S. Alton, Richard G. Bakkala, Gary E. Walters, and Peter T. Munro. December 1988, 31 p.

72. Age determination methods for northwest Atlantic species, edited by Judy Penttila and Louise M. Dery. December 1988, 135 p.

73. Marine flora and fauna of the Eastern United States. Mollusca: Cephalopoda, by Michael Vecchione, Clyde F.E. Roper, and Michael J. Sweeney. February 1989, 23 p.

74. Proximate composition and fatty acid and cholesterol content of 22 species of northwest Atlantic finfish, by Judith Krzynowek, Jenny Murphy, Richard S. Maney, and Laurie J. Panunzio. May 1989, 35 p.

75. Codend selection of winter flounder *Pseudopleuronectes americanus*, by David G. Simpson. March 1989, 10 p.

76. Analysis of fish diversion efficiency and survivorship in the fish return system at San Onofre Nuclear Generating Station, by Milton S. Love, Meenu Sandhu, Jeffrey Stein, Kevin T. Herbinson, Robert H. Moore, Michael Mullin, and John S. Stephens, Jr. April 1989, 16 p.

77. Illustrated key to the genera of free-living marine nematodes of the order Enoplida, by Edwin J. Keppner and Armen C. Tarjan. July 1989, 26 p.

78. Survey of fishes and water properties of south San Francisco Bay, California, 1973–82, by Donald E. Pearson. August 1989, 21 p.

79. Species composition, distribution, and relative abundance of fishes in the coastal habitat off the southeastern United States, by Charles A. Wenner and George R. Sedberry. July 1989, 49 p.

80. Laboratory guide to early life history stages of northeast Pacific fishes, by Ann C. Matarese, Arthur W. Kendall, Jr., Deborah M. Blood, and Beverly M. Vinter. October 1989, 651 p.

NMFS Honolulu Library
00000000004778

NOAA Technical Report NMFS 84

Results of Abundance Surveys of Juvenile Atlantic and Gulf Menhaden, *Brevoortia tyrannus* and *B. patronus*

Dean W. Ahrenholz
James F. Guthrie
Charles W. Krouse

December 1989

U.S. DEPARTMENT OF COMMERCE
Robert Mosbacher, Secretary
National Oceanic and Atmospheric Administration
John A. Knauss, Under Secretary for Oceans and Atmosphere
National Marine Fisheries Service
James Brennan, Assistant Administrator for Fisheries

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Acknowledgments **iv**

Abstract **1**

Introduction **1**

Exploratory-developmental period **2**

Trial-implementation period **3**

Verification analysis **5**
- Estimation of relative abundance **5**
- Correlation tests of survey means with year-class strength estimates **7**
- Trends in regional abundance **8**

Critical conditions and assumptions **8**
- Inherent relationship between juveniles and recruits **8**
- Representative nature of trawl catches **9**
- Timing of the survey **10**
- Proportional representation of regional stream densities **11**
- Uniform format of survey **11**
- Suitability of independent estimates of year-class strength **12**

General discussion and conclusions **12**

Citations **13**

Appendix **14**

Acknowledgments

We are indebted to numerous individuals who made various contributions to the continuation of the surveys over a long period of time. Many helpful analytical suggestions were provided by Alexander J. Chester, David R. Colby, and Rodney L. Reish. During various stages of report preparation, excellent reviews were given by Alexander J. Chester, Sheryan P. Epperly, Charles S. Manooch, III, Walter R. Nelson, Allyn B. Powell, John V. Merriner, and Douglas S. Vaughan. The manuscript has benefitted from discussions with William R. Turner and George N. Johnson, as well as from comments by anonymous reviewers. We thank Herbert R. Gordy for figure preparation, Beverly W. Harvey for word processing, and Curtis W. Lewis for the two-boat trawling illustration.

Results of Abundance Surveys of Juvenile Atlantic and Gulf Menhaden, *Brevoortia tyrannus* and *B. patronus*

DEAN W. AHRENHOLZ
JAMES F. GUTHRIE
CHARLES W. KROUSE
Beaufort Laboratory
Southeast Fisheries Center
National Marine Fisheries Service, NOAA
Beaufort, North Carolina 28516-9722

ABSTRACT

The estuarine populations of juvenile Atlantic and gulf menhaden (*Brevoortia tyrannus* and *B. patronus*) were sampled during two-boat, surface-trawl, abundance surveys extensively conducted in the 1970s. Juvenile Atlantic menhaden were sampled in 39 estuarine streams along the U.S. Atlantic coast from northern Florida into Massachusetts. Juvenile gulf menhaden were sampled in 29 estuarine streams along the Gulf of Mexico from southeast Texas into western Florida. A stratified, two-stage, cluster sampling design was used. Annual estimates of relative juvenile abundance for each species of menhaden were obtained from catch-effort data from the surveys. There were no significant correlations, for either species, between the relative juvenile abundance estimates and fishery-dependent estimates of year-class strength. From 1972 to 1975, the relative abundance of juvenile Atlantic menhaden in north Atlantic estuaries decreased to near zero.

Introduction

Fishery managers entrusted with determining population status, or more critically, the permissible level of harvest for fisheries with variable levels of recruitment, would be aided considerably by estimates of prerecruit year-class strength. With an additional knowledge of recruitment patterns, recommendations could ultimately be made that would maximize yield while reducing risk for a particular fishery resource. Biologists can use a variety of fishery-independent sampling techniques, such as sampling-gear catch-effort indices, hydroacoustic and sonar surveys, or mark-recapture programs, for obtaining estimates of prerecruit year-class strength.

The U.S. Fish and Wildlife Service, Bureau of Commercial Fisheries,[1] expanded their menhaden research program in 1955 to determine whether seasonal fluctuations in commercial purse-seine landings of Atlantic menhaden (*Brevoortia tyrannus*) could be forecasted. This research received additional impetus in the 1960s as landings by the commercial purse-seine fleet steadily declined from the record highs of the mid-1950s (Ahrenholz et al. 1987). One approach used to pursue this objective was to estimate the relative abundance of prerecruit juvenile (young-of-the-year) Atlantic menhaden (Henry 1971, Turner 1973). In addition to the juvenile Atlantic menhaden sampling, which began in 1956, sampling for juvenile gulf menhaden (*B. patronus*) was started in 1964. In the early 1970s, sampling activities culminated in extensive Atlantic and gulf juvenile-abundance trawl surveys that were conducted through 1978. After 1978, the surveys were discontinued because preliminary analyses revealed that the objective of forecasting year-class strength with the survey data could probably not be attained (Ahrenholz et al. 1979). Results of subsequent analyses (presented here) with completed data sets confirmed the earlier finding.

The objectives of this report are to 1) present aspects of estuarine sampling that should be considered when attempting to estimate juvenile abundance, especially of a marine migrant species; 2) demonstrate the need, as well as the large time requirement for verifying juvenile survey results; and 3) document these menhaden surveys, since a large amount of ecological and biological information on menhaden were obtained from these studies.

In this report we first discuss the chronology and salient aspects of what we have termed the exploratory-developmental period of juvenile menhaden abundance sampling. During this period, biologists examined the results for several sampling techniques. We then discuss the trial-implementation period, when extensive, systematic trawl surveys were conducted. We then conduct verification analyses of the survey data. Finally, we examine some initial assumptions relating to geographical and temporal aspects of the sampling design, and suggest some additional directions for the enumeration attempts.

[1]Presently the National Marine Fisheries Service, NOAA, U.S. Dept. of Commerce.



Figure 1
Juvenile gulf menhaden survey streams, 1971–78.

Exploratory-developmental period

Research emphasis for estimating year-class strength focused on the obligatory estuarine phase of the Atlantic menhaden's life history, following suggestions made by Massmann (1953) and Massmann et al. (1954). Because menhaden occupy relatively small estuarine streams during their late-larval and juvenile stages, each developing year-class appeared to be readily available for sampling. Furthermore, earlier workers believed this would result in more accurate estimates than those from larvae that occur in the ocean (Turner 1973).

Sampling began in 1956 and consisted of mark-recapture studies conducted in an estuarine tributary of Indian River, Delaware (Pacheco and Grant 1966). The program expanded into other areas and employed additional sampling techniques. By 1961, abundances of juvenile menhaden inhabiting four coastal estuarine streams (Broad Creek, North Carolina; Felgate Creek, Virginia; White Creek, Delaware; and Childs River, Massachusetts) were estimated using mark-recapture methods and catch-effort data from beach seining and surface trawling. These sampling methods were used in an increasing number of estuaries along the Atlantic coast through 1966. Although the results of all three were similar, Turner (1973) concluded that the surface trawl was the most expedient and widely applicable method. Trawling became the only estuarine sampling method retained. The number of streams sampled by surface trawl increased from 10 in 1962 to 65 in 1968. Trawl sampling for juvenile Atlantic menhaden was not conducted during 1970 or 1971.



Figure 2
Juvenile Atlantic menhaden survey streams, 1972–78.

In 1964, the program was expanded to estimate the relative abundance of juvenile gulf menhaden along the U.S. coast of the Gulf of Mexico (Turner et al. 1974). Since the general life history of this species is similar to that of the Atlantic menhaden, the same basic strategy of sampling was employed. Surface trawling was the only sampling method used.

In addition to the sampling methods discussed, an aerial survey was initiated in 1962 on both the Atlantic and Gulf coasts to estimate the fish-school surface area of juveniles migrating from coastal estuarine streams into coastal waters. Plagued by several operational and interpretational problems, the aerial surveys were discontinued after 1971 (Ahrenholz et al. 1979).

Trial-implementation period

Surface-trawl surveys were conducted with nearly constant sampling effort from 1971 along the U.S. Gulf of Mexico coast and from 1972 along the U.S. Atlantic coast, through



Figure 3
Operation of surface trawl for juvenile Atlantic and gulf menhaden surveys.
(Drawing courtesy of Curtis Lewis, Beaufort Lab., Natl. Mar. Fish. Serv.)

1978. These surveys represented an application of knowledge acquired during the earlier feasibility sampling. Data from the 1970s were comparable and were used to estimate year-class strength and, thus, the efficacy of the survey design.

Estuarine streams selected as sampling areas for each coast appeared to be representative of other streams in the respective surrounding areas. These streams were, for the most part, drawn from the larger set of estuarine streams that had been sampled during the earlier exploratory-developmental period of the program. Selection of a stream during the earlier portion of the program, or the final inclusion into the survey itself, was not purely random. Selection required that each stream be accessible with a launching ramp for trailered boats. Secondly, the streams would have at one time or another been shown to contain menhaden. Estuarine locations ranged from eastern Texas into western Florida on the Gulf of Mexico (Fig. 1), and from northern Florida into Massachusetts along the Atlantic coast (Fig. 2).

Each stream was sampled by a series of tows evenly distributed along the length of the stream. Sites generally ranged from an upstream limit, where the stream was just large enough to deploy sampling gear, to the stream mouth. Each stream would normally have a salinity gradient along its length, which usually ranged from somewhere near 0‰ to 10 or 20‰. The actual number of tow sites per stream varied, ranging on the Gulf coast from 3 to 11, and on the Atlantic from 3 to 8.

Because of the large number of estuaries and vast area involved, each estuary was sampled only once during a single annual survey. The Gulf of Mexico estuaries were surveyed in June, Atlantic estuaries from Florida to New Jersey in June and July, and the estuaries north of New Jersey in September.

Since juvenile menhaden are not effectively sampled in daylight hours when the water is relatively clear, those streams which normally had relatively clear water were sampled at night. Night sampling was, however, impractical to adopt for all streams. On the Gulf coast, streams east of the Mississippi River delta were surveyed at night. On the Atlantic coast, several individual streams from Florida to New Jersey, as well as all streams north of New Jersey, were sampled at night.

Each tow site was sampled by a 5-minute surface-trawl haul. A standardized surface trawl was pulled between two outboard-motor boats (Fig. 3). The trawl was modified after one reported by Massmann et al. (1952) (Fig. 4). Variables recorded for each tow site were surface salinity, secchi depth, temperature, and number of juvenile menhaden captured. Relatively small catches were saved in their entirety, while large catches were subsampled. Samples were preserved in formaldehyde solution. The juvenile menhaden were counted and measured (fork length) in the laboratory.



Figure 4
Standard surface trawl used in juvenile Atlantic and gulf menhaden surveys. Head rope 7.5 m, foot rope 6.7 m, length 6.1 m, and height 0.9 m. Constructed of 6 mm bar-length multifilament knotted nylon netting.

Table 1
Planned distribution of sampling effort by region and estuarine stream for juvenile Atlantic menhaden survey, 1972–78.

Region	Number of streams	Number of tows
North Atlantic	8	32
Middle Atlantic	5	27
Chesapeake Bay	10	50
South Atlantic	16	91
Totals	39	200

Table 2
Planned distribution of sampling effort by region and estuarine stream for juvenile gulf menhaden survey, 1971–78.

Region	Number of streams	Number of tows
Texas	5	38
Louisiana	16	111
Eastern gulf	8	45
Totals	29	194

Verification analysis

Catch-per-tow (CPT) values calculated from trawl samples are measures of relative juvenile menhaden abundance, not actual (or absolute) population size. A series of consistently derived, annual estimates of relative juvenile abundance can be functionally compared (e.g., by regression analysis) with a second, independently derived estimate of absolute population size, and thus "calibrated" (see Eberhardt 1978). The estimates of absolute population (year-class) size for this study were derived from commercial-fishery catch statistics. If a significant, positive relationship exists, the results of the survey have been verified, i.e., there is a high probability that the survey is providing the type of information it was designed to provide.

Estimation of relative abundance

Sampling activities can be described by a stratified two-stage cluster sampling design. Each coast was divided into several regions (strata). Individual streams were the primary sampling units, and individual tows were taken as secondary units. For analysis, the Gulf coast was divided into three geographic regions: Texas, Louisiana, and the eastern Gulf (Mississippi, Alabama, and the Gulf coast of Florida) (Fig. 1). The Atlantic coast was divided into four regions: South Atlantic, Chesapeake Bay, middle Atlantic, and north Atlantic (after June and Reintjes 1959) (Fig. 2). The total numbers of streams and tow sites by region planned for the Atlantic and Gulf coast are given in Tables 1 and 2. Weighting factors for regional means were obtained from estimates of relevant habitat area given in the literature (USFWS 1970) (Tables 3 and 4). Estimates of estuarine area were retained as acre units (as opposed to hectares) for convenience in calculation, as each tow of the trawl sampled approximately one surface acre. Hence, each acre of potential estuarine habitat represents a potential secondary sampling unit for purposes of the computations.

The weighted-mean CPT (estimate of relative abundance) and confidence intervals for each coast and year were then calculated using formulae from Cochran (1963) (Appendix). Regardless of size, all fish captured by the trawl were assigned equal weight to estimate the mean. Results for this series of calculations, done with arithmetic (nontransformed) data, are given in Tables 5 and 6 for the Atlantic and Gulf surveys. Results for computations with $\log_e(y+1)$ transformations are presented as antilogarithms minus one in Tables 7 and 8. Since the antilogarithms of the means (minus one) of the transformed data closely approximate geometric means, they are referred to as such here.

Because of the fixed (nonrandom) nature of the particular streams and tow sites, these estimates of relative abundance are biased. This is not a serious problem, however. With respect to within-years, the only real violation of the random assumption is that the tows are arranged more or less systematically along the length of each estuarine stream. This characteristic increases the probability that the entire range of fish densities in a stream is sampled. With respect to among-year comparisons, it was assumed that differences (variance) in CPT associated with stream and local site factors would be minimized among-years by using the same sites.

Table 3

Estimated habitat of Atlantic menhaden in U.S. Atlantic estuarine areas, adjusted for losses to dredge and fill operations, and used to obtain regional weights for juvenile abundance estimates.

Region and state	Acres[1]
North Atlantic	
Massachusetts	29,000
Rhode Island	13,800
Connecticut	18,200
New York	112,700
Total	173,700
Middle Atlantic and Chesapeake Bay[2]	
New Jersey	357,400
Delaware	143,900
Maryland	375,300
Virginia[3]	425,700
Total	1,302,300
South Atlantic	
North Carolina	785,700
South Carolina	265,100
Georgia	124,200
Florida[4]	73,650
Total	1,248,650

[1]USFWS (1970)
[2]Virginia and Maryland acreage was divided between middle Atlantic (0.15) and Chesapeake Bay (0.85).
[3]Middle Atlantic acreage totaled 621,450 and Chesapeake Bay totaled 680,850.
[4]This value represents 10% of all Florida's estuaries and is assumed to represent that portion of the state's estuaries within the coastal range of Atlantic menhaden.

Table 4

Estimated habitat of gulf menhaden in U.S. Gulf of Mexico estuarine areas, adjusted for losses to dredge and fill operations, and used to obtain regional weights for juvenile abundance estimates.

Region	Acres[1]
Texas	260,000
Louisiana	2,011,500
Eastern Gulf	
Mississippi	74,600
Alabama	130,800
Florida[2]	184,125
Total	389,525

[1]USFWS (1970)
[2]25% of Florida acres were considered to be within the range of gulf menhaden.

Table 5

Actual sampling effort, annual stratified arithmetic mean catch/tow, and 95% confidence intervals (CI) for juvenile Atlantic menhaden survey, 1972–78.

Year class	No. of streams	No. of tows	y_{st}	95% CI
1972	39	200	3215.8	<0.0–6738.8
1973	39	200	1578.2	639.8–2516.6
1974	39	200	1417.4	741.4–2093.4
1975	39	200	653.3	392.4– 914.2
1976	39	200	2753.5	830.7–4676.3
1977	36	183	5248.5	951.4–9545.6
1978	39	200	1850.4	442.8–3258.0

Table 6

Actual sampling effort, annual stratified arithmetic mean catch/tow, and 95% confidence intervals (CI) for juvenile gulf menhaden survey, 1971–78.

Year class	No. of streams	No. of tows	y_{st}	95% CI
1971	29	194	10,517.2	5,716.3–15,318.1
1972	29	191	3,448.4	1,790.3– 5,106.5
1973	29	194	7,439.9	2,031.4–12,848.4
1974	27	186	7,363.8	2,752.9–11,974.7
1975	27	183	2,580.8	520.4– 4,641.2
1976	29	191	6,927.1	2,864.5–10,989.7
1977	29	194	3,626.8	1,457.5– 5,796.1
1978	29	194	8,732.9	4,146.1–13,319.7

Table 7

Actual sampling effort, annual stratified geometric mean catch/tow, and 95% confidence intervals (CI) for juvenile Atlantic menhaden survey, 1972–78.

Year class	No. of streams	No. of tows	y_{st}	95% CI
1972	39	200	86.7	28.0–264.7
1973	39	200	88.9	39.8–197.0
1974	39	200	52.7	21.5–127.3
1975	39	200	27.2	11.3– 63.4
1976	39	200	70.8	29.0–171.5
1977	36	183	135.6	28.0–633.3
1978	39	200	59.1	24.6–141.2

Table 8

Actual sampling effort, annual stratified geometric mean catch/tow, and 95% confidence intervals (CI) for juvenile gulf menhaden survey, 1971–78.

Year class	No. of streams	No. of tows	y_{st}	95% CI
1971	29	194	950.1	421.5–2145.2
1972	29	191	212.7	88.3– 510.6
1973	29	194	497.5	167.6–1474.1
1974	27	186	622.6	277.0–1399.1
1975	27	183	245.3	105.4– 564.3
1976	29	191	490.0	237.4–1010.2
1977	29	194	352.3	137.0– 903.5
1978	29	194	407.0	128.3–1287.4



Figure 5

Stratified arithmetic mean catch-per-tow of juvenile Atlantic menhaden in relation to subsequent recruitment at age 0.5.



Figure 6

Stratified geometric mean catch-per-tow of juvenile Atlantic menhaden in relation to subsequent recruitment at age 0.5.



Figure 7

Stratified arithmetic mean catch-per-tow of juvenile gulf menhaden in relation to subsequent recruitment at age 1.0.

Correlation tests of survey means with year-class strength estimates

Virtual population analysis (VPA) year-class size estimates for Atlantic menhaden from the 1972–78 year-classes were obtained from Ahrenholz et al. (1987), and for gulf menhaden from the 1971–78 year-classes from Vaughan (1987). Correlation analyses of the arithmetic and geometric weighted-mean CPT estimates with appropriate recruitment estimates were performed. Resultant correlation coefficients were not significant ($p>0.10$). Any apparent relationships are too weak to be useful for forecasting year-class strength (Figs. 5–8).



Figure 8

Stratified geometric mean catch-per-tow of juven gulf menhaden in relation to subsequent recruitment at e 1.0.



Figure 9

Log_e of the geometric mean catch-per-tow of juvenile Atlantic menhaden in the north Atlantic area, in the southern portion of the south Atlantic area, and in all areas combined.

Trends in regional abundance

Although combined regional estimates of abundance appear to be insensitive to the degree of fluctuations actually observed in year-class strength, some regional trends in reproductive success were documented by the survey. In the north Atlantic area during 1972–75, numbers of juveniles caught per tow fell from a geometric mean of about 200 to near 1, and did not recover during the remaining years of the survey (through 1978) (Fig. 9). Based on field observations associated with other studies, recovery had not occurred through the 1985 year-class.

A similar but less extreme reduction in reproductive success has occurred in the more southerly portion of the south Atlantic area (from Calabash Creek, North Carolina, to northern Florida). Fluctuations in abundance for the remainder of the south Atlantic area and the remaining two regions more or less parallel total abundance estimates.

Critical conditions and assumptions

Although many minor factors can affect the relationship of relative abundance estimates and estimates of year-class strength, a strong correlation would be expected if the following major conditions were met:

1 A strong functional relationship exists between the true number of juveniles alive at the time of the survey and the number that are subsequently recruited into the fishery.

2 Catches per 5-minute trawl sample reflect what is actually in the stream. (A consistent bias is acceptable here.)

3 The annual survey was conducted at a time when juvenile populations in the estuarine streams are reasonably stable, comprising individuals that will compose the bulk of the recruited year-class, and are of a size that makes them vulnerable to the trawl (~35–60 mm).

4 Regional stream densities are a constant proportion of the population in that general area each year. Stream densities could have a degree of bias, higher or lower than surrounding estuarine waters (bays, sounds, and rivers), as long as the bias is consistent among years.

5 The survey is conducted in a uniform manner each year (e.g., same gear, fished the same).

6 A suitable, reasonably accurate (or consistently biased) and precise estimate of recruited year-class strength is used.

Since a poor correlation exists between survey values and independent fishery estimates of abundance, any or all of the above conditions are suspect.

Inherent relationship between juveniles and recruits

A fundamental but apparently tacit assumption of all the juvenile sampling activity was that year-class strength was established by (or was very early into) the juvenile stage. This notion is generally accepted for most fishes (Savoy and Crecco 1988). A significant relationship between the relative abundance estimates and VPA year-class strength estimates could have provided direct evidence for this assumption.


Atlantic Menhaden – Mean Length
at Age Zero of Fish Landed During
4th Quarter of Fall Fishery
r=–.44*
log R/L r=–.48*
RECRUITMENT X 10⁶ OF ATLANTIC MENHADEN
MEAN LENGTH AGE ZERO (mm)

Figure 10

Estimates of year-class recruitment at analytical age 0.5 against mean length of young-of-the-year Atlantic menhaden landed December–January off North Carolina (* significant at 95% level).


Atlantic Menhaden –Mean Length
at Age Zero of Fish Landed During
3rd Quarter of Chesapeake Bay
Fishery
r=–.63**
log R/L r=–.78**
RECRUITMENT X 10⁶ OF ATLANTIC MENHADEN
MEAN LENGTH AGE ZERO (mm)

Figure 11

Estimates of year-class recruitment at analytical age 0.5 against mean length of young-of-the-year Atlantic menhaden landed September–November in Chesapeake Bay (significant at 99% level).**

Even stronger evidence (but not conclusive) would be provided if the relationship was subsequently persistent in future years of sampling. The converse is not necessarily true, however. The surveys were designed to take advantage of this relationship, not to test for it.

A relationship between numbers of Atlantic menhaden juveniles and subsequent numbers of recruits was observed indirectly by examining density-dependent growth. There was a significant inverse relationship between year-class size and the mean size (fork length) in purse-seine landings (Ahrenholz et al. 1979, Reish et al. 1985). This relationship is reviewed here with additional data. The average sizes of age-0 fish in purse-seine landings were inversely correlated with VPA estimates of year-class strength at age 0.5 (Figs. 10, 11). Since fork length is inversely correlated to estimates of year-class size, the numbers of juveniles alive during the estuarine growth period should also be correlated (positively) to estimates of year-class size. No similar relationship was found for gulf menhaden sampled from the commercial fishery (Nelson and Ahrenholz 1986). However, Guillory and Bejarano (1980) found a significant inverse relationship between mean length of juvenile gulf menhaden and commercial catch-per-unit-effort (CPUE) for age-1 fish. It should be noted, however, that their CPUE values (surrogate for year-class strength estimates) were not significantly correlated to Vaughan's (1987) VPA estimates of year-class size.

Representative nature of trawl catches

The accuracy and precision of trawl catch rates for estimating abundance are dependent upon local (site-specific) vulnerability of juvenile menhaden to the surface trawl. The width and depth of the stream where a haul is made influence vulnerability. Frequently, higher catches in any general area are made where the stream is narrow and/or shallow. This may be due to a decrease in potential avenues of escape for fish avoiding the trawl, but could be a real difference between habitat types. While the trawl is an effective means of capturing juvenile menhaden in coastal streams, it is less effective for capturing larger individuals in larger rivers and in bays and sounds.

Water clarity affects vulnerability. Quite simply, the clearer the water, the sooner fish can see the trawl and avoid it. A comparison of day- and night-tow trawl catches was made for some Gulf estuarine streams surveyed in 1971. The natural logarithms of the ratio of paired night and day tows were plotted against the mean secchi disc values measured during the day portion of the sampling (Fig. 12). These data pairs display a birectilinear relationship. When the mean secchi value is less than about 45 cm, the relative magnitude (and rate of change) of the difference between night and day catches is less than when the secchi exceeds 45 cm.



Figure 12

Log_e of night/day catch-per-tow of juvenile gulf menhaden in relation to secchi depth (data fitted by eye).

The size of fish affects its vulnerability to a surface trawl. Small fish can pass through the mesh, and larger fish can more readily avoid the trawl. Kroger et al. (1974) hypothesized that the trawl is best suited for juvenile menhaden 35–60 mm in fork length. Because it is believed that the survey was conducted at a time when most juveniles fall within this size range, size-induced sampling bias was not evaluated. A theoretical treatment of this potential problem is given by Kjelson and Colby (1976).

Any factors which affect the surface schooling behavior of young menhaden also affect the vulnerability of these fish to the trawl, because only the upper portion of the water column is sampled. Effects of temperature, tide, time of day, and wind on surface schooling of menhaden are unknown. Despite these difficulties, comparisons of trawl catches with haul-seine and mark-recapture studies reveal that the trawl is reasonably effective within the bounds described (Turner 1973, Kroger et al. 1974).

Timing of the survey

Survey timing was adopted in the late 1960s from results of earlier mark-recovery tests and catch-effort sampling. Size of fish was a principal criterion, but early workers also felt that the important fall-winter spawned juveniles were vulnerable during the June–July period. The survey was also conducted in September from waters surrounding Long Island, New York, northward to accommodate spring-summer spawned individuals (Kroger et al. 1974).

The suitability of the timing of the survey was evaluated by examining length-frequency distributions of juveniles in the catches. The distributions by stream were examined to determine if some orderly pattern or shape occurred. If an orderly distribution occurred, it was examined to determine if its modal group(s) was (were) within the range of effective vulnerability.



Figure 13

Weighted length-frequency distribution of trawl catches of juvenile Atlantic menhaden in Hancock Creek, North Carolina, 1972–78.

In some years, many estuarine streams whose data displayed orderly length progressions have a skewed length-frequency distribution, in which the left tail (smaller fish) has not yet formed, and the mode is in the late larval-prejuvenile size range. This indicates that larval or prejuvenile Atlantic menhaden may still be immigrating into the estuarine stream at the time of the survey. These small fish are not completely vulnerable to the trawl, as many will pass through the 6 mm bar-mesh tailbag. An example is given for Hancock Creek, North Carolina (Fig. 13). Similar conditions frequently occurred in Virginia and Maryland estuaries. During 1972 and 1978, the sampling season was apparently too early to adequately sample juveniles. During 1973–77 the small fish seemed to have been adequately sampled. Also, there appeared to be an incomplete representation of larger juveniles (>50 mm) in middle Atlantic and Chesapeake Bay estuarine waters. Thus, the survey may have been too late for early-spawned individuals, as these fish may have already emigrated from the sampling area. Similar but less-pronounced problems were also detected with some of the gulf menhaden length-frequencies.



Figure 14

Length-frequency distribution by tow with salinity values for juvenile Atlantic menhaden from the 1977 trawl survey in Hancock Creek, North Carolina. (The 1977 weighted distribution of Figure 10 resulted from these distributions.)



Figure 15

Length-frequency distribution by tow with salinity values for juvenile gulf menhaden from the 1966 trawl sampling activities in Dog River, Alabama.

Proportional representation of regional stream densities

Proportional representation of fish densities in regional streams in the survey was evaluated using length-frequency distributions. It was assumed that a near-constant proportional cross-section of the juvenile population was not sampled each year if patterns of length-frequency distributions were not similar within streams among years, when patterns were differentially mixed among streams within years.

Some survey design problems can be attributed to early assumptions on life history and the concept of "primary nursery area." The hypothesis, developed early in the studies, was that young (late-larval and prejuvenile) menhaden migrate upstream into small estuarine streams, where near 0‰ salinity occurs, and then progressively move downstream as they grow. This hypothesis was supported by observed length-frequencies in a large portion of the streams sampled, as juveniles appeared to be stratified by size along the salinity gradient (Figs. 14 and 15). Not infrequently, however, similar size distributions were observed when no apparent salinity gradient existed (Fig. 16).

Implicit to a survey design based on streams selected for the occurrence of a salinity gradient is the assumption that a population of fish once immigrated into a stream will remain in that stream until emigration in late summer or fall. Often, however, length-frequency distributions have missing portions, i.e., the right, left, or middle of an expected size-frequency distribution was not represented in the catches in one stream but was present in another stream during the same year. Thus, it appears that there are varying degrees of interchange between survey streams and the larger estuarine system to which they are tributary. The interchange appears to be especially common in the Chesapeake Bay system.

Uniform format of survey

Considerable care was exercised during the 1971–78 Gulf of Mexico and 1972–78 Atlantic surveys to keep all aspects of the survey as constant as possible. Some minor problems were created when a stream could not be adequately sampled due to periodic plant growth such as water hyacinths (*Eichornia crassipes*) or duck weed (*Lemna* sp.) as well as by irregular boat or barge traffic (note differences in sampling effort between Tables 1 and 2, and 5–8).



Figure 16

Length-frequency distribution by tow for juvenile gulf menhaden from the 1975 trawl survey in Dog River, Alabama (all salinity measurements were 0‰).

Suitability of independent estimates of year-class strength

Several potentially comparative estimates of year-class strength can be obtained from commercial purse-seine landing statistics and port samples: Landings in biomass in a fixed number of subsequent years, landings in numbers at a fixed age for a given year-class, catch-per-unit-effort for a given age, or VPA or cohort (backward sequential) analysis which estimates number of recruits at a fixed age for a given year-class. Each has advantages and disadvantages.

Estimates of landings in biomass, which during the time of the survey were dominated by age-2 fish on the Atlantic and age-1 fish in the Gulf of Mexico, are readily obtainable and are the most accurate. Strong correlations with survey means would depend on constant growth rates among age-classes, as well as each age group comprising a near-constant percentage of the landings among years. Also, the rate of fishing would have to be nearly constant. These conditions are not well met, so more adjustments would be needed.

To use age-specific landings in numbers as a measure of year-class strength, the rate of fishing on the age group would have to be (nearly) constant from year to year. This condition is only marginally met over short spans of years, so again adjustments need to be made. Sampling errors about estimates for the numbers-at-age landed appear to be at acceptable levels (see Chester 1984, Chester and Waters 1985).

CPUE estimates in some fisheries may be useful as a measure of abundance, if the interacting effects of other age groups can be separated. CPUE is usually inappropriate for use with the commercial menhaden purse-seine fishery. For CPUE to reflect true abundance, the catchability coefficient must be (nearly) constant. It is, however, inversely related to population size for both the Atlantic menhaden (Schaaf 1975) and gulf menhaden (Nelson and Ahrenholz 1986) fisheries.

VPA estimates are the closest to being true estimates of year-class strength. This analysis provides an estimate of number of recruits for a fixed time-period before significant fishing has taken place on the year-class. VPA for Atlantic menhaden should include age-0.5 fish, as large catches of estuarine-emigrating juveniles are frequently made. This is not the case for gulf menhaden, where age-1 is an appropriate recruitment age. A preliminary VPA estimate can be obtained only for the Atlantic fishery after landing and port-sampling data are processed for the complete fishing season on 3-year-old fish of a given year-class. The VPA estimates appear to be stable after the fifth year for the Atlantic and the third year for the Gulf fishery. Because of these constraints, a preliminary correlation analysis of the first 3 years of a survey would not have been possible for either fishery until about 7 years after the survey was initiated.

General discussion and conclusions

Although substantial biogeographical and life-history information for these two species of menhaden resulted from juvenile surveys, the original objective of the sampling was not achieved, i.e., the ability to forecast year-class strength. It appears that the survey suffered from assumptions that were not met.

Following the preliminary life-history studies and technique developments of the 1950s and 1960s, the survey was initiated and the format was "locked in." Since success would depend on the survey being conducted in a constant fashion, any new information pertinent to the survey design could not be implemented. Quite simply, a substantial investment had already been made in research funds and time. Initial assumptions could not be adequately evaluated as to whether or not the survey was effective until a sufficient number of years had passed. Continued adjustments would require additional years of data for testing, since the historic database would be incomplete or noncomparable.

Once sufficient data were in hand, the straightforward comparison of weighted mean CPT, using weighting factors derived independently from survey and catch data, with what are believed to be analytically the best available estimates of year-class strength, is fundamental. The lack of a reasonably strong (and significant) relationship is a logical and prudent stopping point for a survey of this type. A risk is present, however, as some analytical fault may lie with estimates of ages, numbers landed, and other factors associated

with purse-seine, fishery-dependent measures of year-class strength.

The results of an extensive number of analyses were not reported here. These included multiple regressions, various weighting schemes, a number of derived measures of abundance, and stream or regional combinations in correlation analyses with a variety of types of fishery-derived measures of year-class strength. Without a preconceived hypothesis of cause and effect for variables selected in these analyses, approaches of this type are simply trial-and-error searches, and will no doubt result in one or more apparent significant relationships due simply to chance. The risk here, of course, is that one of the potential trial-and-error solutions is correct.

Development of a new survey format should be undertaken to incorporate new information from other studies and types of analyses that were not available or apparent during the inception of the discontinued survey. Emphasis should be placed on the geographic timing of the survey relative to parental spawning in the region, and on obtaining more useful sampling-stratification criteria.

The stratification criteria used here were geographic. These criteria apparently did not relate that strongly to actual differences in juvenile fish density. Significant reductions in the variance about the stratified mean CPT were not gained over a single-stage model (tow being the primary sampling unit), or a purely random sampling model.

Some promise with respect to an alternate approach is offered by the density-dependent growth relationship. While the observed correlations are not extremely strong, at least for forecasting purposes, they are encouraging considering the degrees of bias associated with length parameters derived from samples of purse-seine landings. The purse-seine catches are not systematic relative to time of year, and tend to be selective towards larger individuals. This tends to distort the apparent relationship on the left-hand portion of Figures 10 and 11, particularly for fishery data from North Carolina (Fig. 10).

Because of the selective size bias and attempts to reduce commercial catches of younger-age Atlantic menhaden (ASMFC 1981), no attempt was made to refine this relationship. Estimates of mean length from field sampling, which would parallel relative abundance sampling, may be a plausible approach. The discussions presented earlier on length-frequencies indicate that this aspect of sampling will also require substantial refinement. If this approach proves useful, a verification of the technique may be possible using data from historic surveys, once more in-depth analytical explanations of the observed length-frequency distributions are obtained.

Citations

Ahrenholz, D.W., J.F. Guthrie, and C.W. Krouse
1979 Evaluation of Atlantic and gulf menhaden juvenile abundance surveys. Unpubl. manuscr., Southeast Fish. Cent. Beaufort Lab., Natl. Mar. Fish. Serv., NOAA, Beaufort, NC 28516, 41 p.

Ahrenholz, D.W., W.R. Nelson, and S.P. Epperly
1987 Population and fishery characteristics of Atlantic menhaden, *Brevoortia tyrannus*. Fish Bull., U.S. 85:569–600.

ASMFC
1981 Fishery management plan for Atlantic menhaden. Fish. Manage. Rep. 2, Atlantic States Marine Fisheries Commission, 1400–16th St. N.W., Wash., D.C. 20036, 134 p.

Chester, A.J.
1984 Sampling statistics in the Atlantic menhaden fishery. NOAA Tech. Rep. NMFS 9, Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 16 p.

Chester, A.J., and J.R. Waters
1985 Two-stage sampling for age distribution in the Atlantic menhaden fishery, with comments of optimal survey design. N. Am. J. Fish. Manage. 5:449–456.

Cochran, W.G.
1963 Sampling techniques. Wiley, NY, 413 p.

Eberhardt, L.L.
1978 Appraising variability in population studies. J. Wildl. Manage. 42:207–238.

Guillory, V., and R. Bejarano
1980 Evaluation of juvenile menhaden abundance data for prediction of commercial harvest. Proc. Annu. Conf. Southeast Assoc. Fish Wildl. Agencies 34:193–203.

Henry, K.A.
1971 Atlantic menhaden (*Brevoortia tyrannus*) resource and fishery—analysis of decline. NOAA Tech. Rep. NMFS SSRF 642, Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 32 p.

June, F.C., and J.W. Reintjes
1959 Age and size composition of the menhaden catch along the Atlantic Coast of the United States, 1952–55; with a brief review of the commercial fishery. U.S. Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 317, 65 p.

Kjelson, M.A., and D.R. Colby
1976 The evaluation and use of gear efficiencies in the estimation of estuarine fish abundance. *In* Wiley, M. (ed.), Estuarine processes, vol. 2, p. 416–424. Acad. Press, NY.

Kroger, R.L., A.L. Pacheco, J.F. Guthrie, and L.R. Whitfield
1974 Development and present status of survey methods for estimating relative abundance of juvenile menhaden. Unpubl. manuscr., Southeast Fish. Cent. Beaufort Lab., Natl. Mar. Fish. Serv., NOAA, Beaufort, NC 28516, 36 p.

Massmann, W.H.
1953 Relative abundance of young fishes in Virginia estuaries. Trans. N. Am. Wildl. Nat. Resour. Conf. 18:439–446.

Massmann, W.H., E.C. Ladd, and H.N. McCutcheon
1952 A surface trawl for sampling young fishes in tidal rivers. Trans. N. Am. Wildl. Nat. Resour. Conf. 17:386–392.
1954 Postlarvae and young of the menhaden (*Brevoortia tyrannus*) in brackish and fresh waters of Virginia. Copeia 1954:19–23.

Nelson, W.R., and D.W. Ahrenholz
1986 Population and fishery characteristics of gulf menhaden, *Brevoortia patronus*. Fish Bull., U.S. 84:311–325.

Pacheco, A.L. and G.C. Grant
1966 Part II — Occurrence, abundance, and growth of young menhaden in a tributary creek of Indian River, Delaware, 1957. Unpubl. manuscr., Southeast Fish. Cent. Beaufort Lab., Natl. Mar. Fish. Serv., NOAA, Beaufort, NC 28516, 30 p.

Reish, R.L., R.B. Deriso, D. Ruppert, and R.J. Carroll
1985 An investigation of the population dynamics of Atlantic menhaden (*Brevoortia tyrannus*). Can. J. Fish. Aquat. Sci. 42:147–157.

Savoy, T.F., and V.A. Crecco
1988 The timing and significance of density-dependent and density-independent mortality of American shad, *Alosa sapidissima*. Fish Bull., U.S. 86:467–482.

Schaaf, W.E.
1975 Fish population models: potential and actual links to ecological models. *In* Russell, C.S. (ed.), Ecological modeling in a resource management framework, p. 211–239. Resources for the Future, Wash., D.C.

Turner, W.R.
1973 Estimating year-class strength of juvenile menhaden. *In* Pacheco, A.L. (ed.), Proceedings of a workshop on egg, larval, and juvenile stages of fish in Atlantic coast estuaries, p. 37–47. Tech. Publ. 1, Mid Atl. Coastal Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Highlands, NJ 07732.

Turner, W.R., G.N. Johnson, and H.R. Gordy
1974 Compendium of juvenile menhaden surveys in coastal streams of the northern Gulf of Mexico. Data Rep. 89, Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 189 p. (microfiche).

USFWS
1970 National estuary study, vol. 5, p. 82–83. U.S. Fish Wildl. Serv., Wash., D.C. 20242.

Vaughan, D.S.
1987 A stock assessment of the gulf menhaden, *Brevoortia patronus*, fishery. NOAA Tech. Rep. NMFS 58, Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 18 p.

Appendix

Estimates of weighted mean catch-per-tow were obtained from the following stratified two-stage cluster sampling design. Equations are from Cochran (1963).

Given:

L = number of regions (geographical strata),
N_h = number of potential streams in region h,
n_h = number of streams sampled in region h,
$\bar{M}_h$ = mean stream size (of those sampled) in region h (equivalent to mean number of potential tows),
M_{hi} = size of stream i, region h,
m_{hi} = number of tows in stream i, region h,
f_{1h} = finite population correction for streams by region (n_h/N_h),
f_{2hi} = finite population correction for tows by stream (m_{hi}/M_{hi}),
y_{hij} = catch-per-tow j, in stream i, region h,
$\bar{y}_{hi}$ = mean catch-per-tow in stream i, region h,
$\bar{\bar{y}}_{R,h}$ = weighted mean catch-per-tow for region h,
$\bar{\bar{y}}_{st}$ = stratified mean catch-per-tow for entire coast,
W_h = regional weight $(N_h\bar{M}_h / \sum_{h=1}^{L} N_h\bar{M}_h)$,
$v(\ )$ = estimate of the variance of an estimate ( ), and
s_{2hi}^2 = estimate of variance of catch-per-tow for stream i, region h.

Estimates of means and variances were obtained from:

$$\bar{y}_{hi} = \frac{\sum_{j=1}^{m_{hi}} y_{hij}}{m_{hi}} \tag{1}$$

$$\bar{\bar{y}}_{R,h} = \frac{\sum_{i=1}^{n_h} \bar{y}_{hi} M_{hi}}{\sum_{i=1}^{n_h} M_{hi}} \tag{2}$$

$$\bar{\bar{y}}_{st} = \sum_{h=1}^{L} W_h \bar{\bar{y}}_{R,h} \tag{3}$$

$$s_{2hi}^2 = \frac{\sum_{j=1}^{m_{hi}} (y_{hij} - \bar{y}_{hi})^2}{m_{hi} - 1} \tag{4}$$

$$v(\bar{\bar{y}}_{R,h}) = \frac{1 - f_{1h}}{n_h \bar{M}_h^2} \cdot \frac{\sum_{i=1}^{n_h} M_{hi}^2 (\bar{y}_{hi} - \bar{\bar{y}}_{R,h})^2}{n_h - 1} + \frac{f_{1h}}{n_h^2 \bar{M}_h^2} \cdot \sum_{i=1}^{n_h} \frac{M_{hi}^2 (1 - f_{2hi}) s_{2hi}^2}{m_{hi}}, \tag{5}$$

$$v(\bar{\bar{y}}_{st}) = \sum_{h=1}^{L} W_h^2 \, v(\bar{\bar{y}}_{R,h}). \tag{6}$$

Confidence intervals (CI) were estimated as follows using the appropriate value of Students t:

$$95\%\ \text{CI} = \bar{\bar{y}}_{st} \pm t\,[v(\bar{\bar{y}}_{st})]^{1/2}. \tag{7}$$

NOAA TECHNICAL REPORT NMFS

Guidelines for Contributors

FORM OF MANUSCRIPT

Authors are encouraged to retain manuscripts on word-processing storage media (diskettes, floppy disks) and submit a double-spaced hardcopy run from the storage media. The document should be in the following sequence: Title page (incl. authors' full names and complete mailing addresses), Contents, Acknowledgments, Abstract, Text, Citations, Text footnotes, Appendices, Tables, Figure legends, and Figures.

Disks Submit disks as MS-DOS "print" or "non-document" files (often called "ASCII files"). If a disk cannot be converted to an ASCII file, the author should indicate on the disk the source computer and software language (e.g., Word Star, Word Perfect) along with the file name. Either 5¼-inch or 3½-inch disks from IBM-compatible or Apple/MacIntosh systems (non-graphics only) can be submitted, double-sided/double-density or high-density, limiting each file to 300 kilobytes. All 8-inch word-processing disks (e.g., Wang or NBI) must be converted onto 5¼- or 3½-inch MS-DOS print disks.

CONTENTS OF MANUSCRIPT

Acknowledgments Gather all acknowledgments into a brief statement at the beginning of the text. Give credit only for exceptional contributions and not to those whose contributions are part of their normal duties.

Abstract Not to exceed one double-spaced typed page. Should include a sentence or two explaining to the general reader why the research was undertaken and why the results should be viewed as important. Abstract should convey the main point of the paper and outline the results or conclusions. No footnotes or references.

Text A brief introduction should portray the broad significance of the paper. The entire text should be intelligible to readers from different disciplines. All technical terms should be defined. Follow the *U.S. Government Printing Office Style Manual*, 1984 ed., and the *CBE Style Manual*, 5th ed. Fishery nomenclature should follow the American Fisheries Society Special Publication 12, *A List of Common and Scientific Names of Fishes from the United States and Canada*, 4th ed.

Abbreviations and symbols Define all abbreviations, acronyms, and symbols in text, equations, and formulae. Abbreviate units of measure only when used with numerals or in tables and figures to conserve space.

Measurements should be expressed in metric units. Other equivalent units may be given in parentheses.

Text footnotes should be numbered in Arabic numerals and typed on a separate sheet from the text. Footnotes are not used for reference material or personal communications, but rather to explain or define terms in the text and for contribution numbers.

Personal communications are noted parenthetically in the text (full name, affiliation, brief address including zip code, month and year).

Citations All written sources should be listed in the Citations section, including unpublished and processed material. In text, cite as Smith and Jones (1977) or (Smith and Jones 1977); if more than one citation, list chronologically (Smith 1936, Jones 1975, Doe 1977). All sources cited in the text should be listed alphabetically by the senior authors' surname under the heading CITATIONS. Abbreviations of periodicals and serials should conform to the Biosis Data Base™. Indicate whether sources are in a language other than English. For informal literature, include address of author or publisher. Authors are responsible for the accuracy of all citations.

Tables should supplement, not duplicate, the text. Each table should be numbered and cited consecutively, with headings short but amply descriptive so that the reader need not refer to the text. For values less than one, zeros should precede all decimal points. Indicate units of measure in column headings; do not deviate from the unit of measure within a column. Table footnotes should be noted consecutively in Roman letters across the page from left to right and then down. Since all tables are typeset, they need not be submitted camera-ready.

Figures Photographs and line drawings should be of professional quality—clear and concise—and reducible to 42 picas for full-page width or to 20 picas for a single-column width, and a maximum of 57 picas high. All graphic elements in illustrations must be proportioned to insure legibility when reduced to fit the page format. Line weight and lettering should be sharp and even. Lettering should be upper and lower case, and vertical lettering should be avoided whenever possible (except for vertical, y, axis). Zeros should precede all decimal points for values less than one. Reproductions of line artwork are accepted in the form of high-quality photographic prints from negatives or photomechanical transfer (PMT). Halftones should be sharply focused with good contrast. Micron rules should be inserted on electron micrographs, even when magnification is included in the figure legend. There should be clear distinction between identifying letters (press-on or overlays) and background of photograph. Label each figure in pencil on the back. Send only xerox copies of figures to the Scientific Editor; originals or photographic prints will be requested later when the manuscript is accepted for publication.

Copies of published reports For single monographs, the senior author and his/her organization each receive 50 copies free-of-charge. For collections of papers, the senior author and his/her organization each receive 50 separates and one cover free-of-charge. Additional copies may be purchased in lots of 100.

Copyright Government publications are in the public domain, i.e., they are not protected by copyright.

SUBMISSION OF MANUSCRIPTS

Send original hardcopy and two duplicated copies to:

Dr. Linda L. Jones, Scientific Editor
National Marine Mammal Laboratory, F/AKC3
National Marine Fisheries Service, NOAA
7600 Sand Point Way N.E.
Seattle, WA 98115-0070

Indicate whether word processing storage media are available, but do not forward disks until requested by Scientific Publications Office.

Questions regarding editorial policy and procedures or disk specifications may be forwarded to:

Nancy Peacock, Managing Editor
NOAA Technical Reports NMFS
National Marine Fisheries Service, NOAA
Scientific Publications Office, F/NWR1
7600 Sand Point Way N.E.
Seattle, WA 98115-0070

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC
ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS OFFICE
BIN C15700
SEATTLE, WA 98115

OFFICIAL BUSINESS

Penalty for Private Use, $300

BULK RATE
POSTAGE & FEES PAID
U.S. Department of Commerce
Permit No. G-19

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 13, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; predictions and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161